EXOTIC PEST FACT SHEET 2

Vegetable Leaf Miners (Liriomyza spp.)



What are they?

Vegetable leaf miners (including *Liriomyza trifolii, L. strigata, L. sativae, L. huidobrensis* and *L. pusilla*) are small flies whose larvae feed internally on plant tissue, particularly the leaf. Feeding by the larvae disrupts photosynthesis and reduces plant quality and yield.

What are the main hosts?

Leaf miners have a very wide host range, including many economically important grains and vegetables. The main processed vegetable crops are beetroot, carrots, beans and peas.

What do they look like?

The black flies are just visible (1-2.5 mm long) and often have yellow spots on the head and thorax (Fig 1). Leaf mines caused by larval feeding are usually white coloured with dampened black and dried brown areas (Fig 2, 3). These mines are typically serpentine or irregularly shaped, and increase in size as the larvae mature. Damage to the plant is caused in several ways:

- Leaf stippling resulting from females feeding or laying eggs.
- · Internal mining of the leaf by the larvae.
- Secondary infection by pathogens that enter the leaf mines or punctures.
- Mechanical transmission of viruses.

Why are they an issue?

Many species of leaf miners are unwanted pests for New Zealand. Of particular concern is *L. huidobrensis* (Serpentine leaf miner) which is listed as a priority pest by Biosecurity New Zealand.

What should I look for?

A leaf miner infestation is most likely detected by the presence of mines in leaf tissue. Adult flies and larvae are unlikely to be seen because of their small size.

How do they spread?

Adult flies can spread throughout a crop by flying, but most long distance spread occurs when plant material containing larvae or eggs is transported.

Where are they present?

Leaf miners of concern to New Zealand are generally widespread through Africa, America, Europe, Asia and parts of Oceania. The species *Liriomyza sativae* is present in Australia in north Queensland.

How can I protect my industry?

Check your production sites frequently for the presence of new diseases and unusual symptoms. Make sure you are familiar with common pests and diseases of your industry so you can recognise something different.



Fig 1. Adult *Liriomyza huidobrensis* leaf miner. Image: National Plant Protection Organisation, The Netherlands, Bugwood.org



Fig 2. A common bean leaf showing tracts from leaf miner infestation. Image: Howard F. Schwartz, Colorado State University, Bugwood.org.



Fig 3. Leaf mines in a squash leaf caused by *Liriomyza* leaf miners. Image: University of Florida.

Version 1. March 2020